3 (Sem-1/CBCS) MAT HC2

## 2021 (Held in 2022)

## **MATHEMATICS**

(Honours)

Paper: MAT-HC-1026

(Algebra)

Full Marks: 80

Time: Three hours

## The figures in the margin indicate full marks for the questions.

- 1. Answer the following as directed:  $1 \times 10 = 10$ 
  - (a) Find the polar representation of z = 2i.
  - (b) If x = 0 and y > 0, then what is the value of  $t^*$ ?
  - (c) Write the negation of the statement 'For any integer n,  $n^2 > n$ ' in plain English then formulate the negation using set of context and quantifier.

- (d) Disapprove the statement using counter example:
  - "For any  $x, y \in \mathbb{R}$ ,  $x^2 = y^2$  implies x = y."
- (e) Suppose f is a constant function from X to Y. The inverse image of a subset of Y cannot be
  - (i) an empty set
  - (ii) the whole set X
  - (iii) a non-empty proper subset of X (Choose the correct option)
- (f) Let  $X = Y = \mathbb{R}$ . Let  $A \subseteq X$ ,  $B \subseteq Y$ . Draw the picture for  $A \times B$  where A = [-1,1] and B = [2,3].
- (g) Suppose a system of linear equations in echelon form has a 3 × 5 augmented matrix whose fifth column is a pivot column.

Is the system consistent? Justify.

(h) If a set  $S = \{\vec{v}_1, \vec{v}_2, ...., \vec{v}_p\}$  in  $\mathbb{R}^n$  contains the  $\vec{O}$  vector, is the set linearly independent? Justify.

- (i) If  $A = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix}$   $\vec{x} = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$ , compute  $(A\vec{x})^T$ .
- (j) What is the determinant of an  $n \times n$  elementary matrix E that has been scaled by 7.
- 2. Answer the following questions: 2×5=10
  - (a) If  $z = -2\sqrt{3} 2i$ , find the polar radius and polar argument of z.
  - (b) Is the function  $g: \mathbb{R} \to \mathbb{R}$  given by g(x) = |x-2| one-one and onto? Explain.
  - (c) Let universal set be  $\mathbb{R}$  and index set be  $\mathbb{N}$ . For a natural number n,  $J_n = \left(0, \frac{1}{n}\right)$ .

Identify with justification  $\bigcap_{n\in\mathbb{N}} J_n$ .

(d) Show that T is a linear transformation by finding a matrix that implements the mapping

$$T(x_1, x_2, x_3, x_4) = (0, x_1 + x_2, x_2 + x_3, x_3 + x_4)$$

- (e) A is a 2 × 4 matrix with two pivot positions. Answer the following with justification:
  - (i) Does  $A\vec{x} = \vec{0}$  have a non-trivial solution?
  - (ii) Does  $A\vec{x} = \vec{b}$  have at least one solution for every  $\vec{b}$ ?
- 3. Answer any four questions from the following: 5×4=20
  - (a) Find the polar representation of the complex number

$$z = 1 - \cos a + i \sin a \quad a \in [0, 2\pi).$$

(b) Let A and B be subsets of an universal set U. Prove —

(i) 
$$(A \cap B)^C = A^C \cup B^C$$

(ii) 
$$(A \cup B)^C = A^C \cap B^C$$
 5

- (c) Define bijection. Let  $f: \mathbb{N} \to \mathbb{N}$  be f(m) = m - 1, if m is even f(m) = m + 1, if m is odd. Show f is a bijection and  $f^{-1} = f$ . 1+4=5
- (d) For vectors  $\vec{v}_1, \vec{v}_2, ...., \vec{v}_p \in \mathbb{R}^n$  define span  $\{\vec{v}_1, \vec{v}_2, ...., \vec{v}_p\}$  construct a  $3 \times 3$  matrix A with non-zero elements and a vector  $\vec{b}$  on  $\mathbb{R}^3$  such that  $\vec{b}$  is not in the set spanned by the columns of A. 2+3=5
- (e) Alka-Seltzer contains sodium bicarbonate ( $NaHCO_3$ ) and citric acid ( $H_3C_6H_5O_7$ ). When a tablet is dissolved in water the following reaction produces sodium citrate, water and carbon dioxide:
- $NaHCO_3 + H_3C_6H_5O_7 \rightarrow Na_3C_6H_5O_7 + H_2O + CO_2$ Balance the chemical equation using vector equation approach. 5
  - (f) Prove that an  $n \times n$  matrix A is invertible if and only if A is row equivalent to  $I_n$ , and in this case any sequence of elementary row operations that reduces A to  $I_n$  also transforms  $I_n$  into  $A^{-1}$ .

- 4. Answer any four from the following:  $10 \times 4 = 40$ 
  - (i) Find the cube roots of the number (a) z = 1 + i and represent them in the complex plane. 5
    - (ii) Find the number of ordered pairs (a, b) of real numbers such that  $(a+ib)^{2002}=a-ib.$ 2
    - (iii) If x, y, z be real numbers such that  $\sin x + \sin y + \sin z = 0$ and  $\cos x + \cos y + \cos z = 0$ , prove that  $\sin 2x + \sin 2y + \sin 2z = 0$ and  $\cos 2x + \cos 2y + \cos 2z = 0$ .

- (i) Solve the equation (b)  $z^7 - 2iz^4 - iz^3 - 2 = 0$ . 5
  - Find the inverse of the matrix if (ii) it exists by performing suitable row operations on the augmented matrix [A: I]

$$A = \begin{bmatrix} 1 & 0 & -2 \\ -3 & 1 & 4 \\ 2 & -3 & 4 \end{bmatrix}$$
 5

- (c) (i) If  $f: X \to Y$  be a map and  $B \subseteq Y$ , then prove  $f^{-1}(B^C) = (f^{-1}(B))^C$ .
  - (ii)  $A_n = \left(-\frac{1}{n}, \frac{1}{n}\right)$ , where  $n \in \mathbb{N}$ . Find  $\bigcup_{n \in \mathbb{N}} A_n \text{ and } \bigcap_{n \in \mathbb{N}} A_n.$  2
  - (iii) Let  $f: \mathbb{R} \to \mathbb{R}$  be given by  $f(x) = x^2$ . Find  $f^{-1}(1), f^{-1}(-1), f^{-1}([0, 1])$ .
- (d) (i) State the induction principle and use it to show that for any positive integer  $1+2+3+...+n=\frac{n(n+1)}{2}$ .

4

(ii) Write as an implication 'square of an even number is divisible by 4'.

Then use direct proof to prove it.

7

- (iii) Give proof using contrapositive 'For an integer x if  $x^2 - 6x + 5$  is even, then x is odd'.
- (e) (i) Use the invertible matrix theorem to decide if A is invertible

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 3 & 1 & 2 \\ -5 & -1 & 9 \end{bmatrix}$$
 2

(ii) Compute det A where

$$A = \begin{bmatrix} 2 - 8 & 6 & 8 \\ 3 - 9 & 5 & 10 \\ -3 & 0 & 1 - 2 \\ 1 - 4 & 0 & 6 \end{bmatrix}$$

(iii) What do you mean by equivalence class for an equivalence relation? For the relation  $a \equiv b \mod(5)$  on z, find all the distinct equivalence classes of z. 1+3=4

(f) (i) Solve the system of equations

$$x_1 - 3x_3 = 8$$
$$2x_1 + 2x_2 + 9x_3 = 7$$
$$x_2 + 5x_3 = -2$$

- (ii) Choose h and k such that the system has 4
  - (a) no solution
  - (b) a unique solution
  - (c) many solutions

$$x_1 + hx_2 = 2$$
$$4x_1 + 8x_2 = k$$

- (iii) Write the general solution of  $10x_1 3x_2 2x_3 = 7$  in parametric vector form.
- (g) (i) Prove that the indexed set  $S = \left\{ \vec{v}_1, \vec{v}_2, ...., \vec{v}_p \right\}$  of two or more vectors is linearly dependent if and only if at least one of the vectors in S is a linear combination of the others. In fact, if S is linearly dependent and  $\vec{v}_1 \neq \vec{0}$ , then some  $\vec{v}_j$  (with j > 1) is a linear combination of the preceding vectors  $\vec{v}_1, \vec{v}_2, ..., \vec{v}_{j-1}$ .

(ii) Use Cramer's rule to compute the solutions of the system 3

$$-5x_1 + 3x_2 = 9$$
$$3x_1 - x_2 = -5$$

- (iii) Suppose  $T: \mathbb{R}^5 \to \mathbb{R}^2$ and  $T(\vec{x}) = A\vec{x}$  for some matrix A and each  $\vec{x}$  in  $\mathbb{R}^5$ . How many rows and columns does A have? Justify.
- (h) (i) Let  $T: \mathbb{R}^2 \to \mathbb{R}^2$  be the transformation that rotates each point in  $\mathbb{R}^2$  about the origin through an angle  $\phi$  with the counter-clockwise direction taken as positive. Find the standard matrix for this transformation.

3

(ii) Let T: R<sup>n</sup> → R<sup>m</sup> be a linear transformation.
 Prove that T is one-to-one if and only if the equation T(x̄)=0 has only the trivial solution.

(iii) Find the area of the parallelogram whose vertices are (0, -2), (6, -1), (-3, 1) and (3, 2).