#### 3 (Sem-5/CBCS) MAT HE 4/5/6

#### 2021

(Held in 2022)

#### MATHEMATICS

(Honours Elective)

### Answer the Questions from any one Option.

#### OPTION-A

Paper: MAT-HE-5046

(Linear Programming)

#### DSE(H)-2

Full Marks: 80

Time: Three hours

# The figures in the margin indicate full marks for the questions.

- 1. Answer the following as directed:
  - $1 \times 10 = 10$
  - (a) A basic feasible solution whose variables are.
    - (i) degenerate
    - (ii) nondegenerate

Contd.

- (iii) non-negative
- (iv) None of the above (Choose the correct answer)
- (b) The inequality constraints of an LPP can be converted into equation by introducing
  - (i) negative variables
  - (ii) non-degenerate B.F.
  - (iii) slack and surplus variables
  - (iv) None of the above (Choose the correct answer)
- (c) A solution of an LPP, which optimize the objective function is called
  - (i) basic solution
  - (ii) basic feasible solution
  - (iii) optimal solution
  - (iv) None of the above (Choose the correct answer)
- (d) What is artificial variable of an LPP?
- (e) Write the equation of line segment in  $\mathbb{R}^n$ .
- (f) Define dual of a given LPP.

- (g) What is pure strategy of game theory?
- (h) Is region of feasible solution to an LPP constitute a convex set?
- (i) Is every convex set in  $\mathbb{R}^n$  a convex polyhedron also?
- (j) Is every boundary point an extreme point of a convex set?
- 2. Answer the following questions:  $2 \times 5 = 10$ 
  - (a) Show that the feasible solution  $x_1 = 1, x_2 = 0, x_3 = 1, z = 6$  to the system  $\min Z = 2x_1 + 3x_2 + 4x_3$  s.t.  $x_1 + x_2 + x_3 = 2$   $x_1 x_2 + x_3 = 2$  ,  $x_i \ge 0$  is not basic.
  - (b) A hyperplane is given by the equation  $3x_1 + 2x_2 + 4x_3 + 7x_4 = 8$ Find in which half space do the point (-6,1,7,2) lie.
  - (c) Find extreme points if any of the set  $S = \{(x,y): |x| \le 1, |y| \le 1\}$
  - (d) Show by an example that the union of two convex sets is not necessarily a convex set.

(e) If 
$$x_1 = 2$$
,  $x_2 = 3$ ,  $x_3 = 1$  a BFS of the LPP max  $Z = x_1 + 2x_2 + 4x_3$   
s.t.  $2x_1 + x_2 + 4x_3 = 11$   
 $3x_1 + x_2 + 5x_3 = 14$   
 $x_1, x_2, x_3 \ge 0$ ? Explain.

- 3. Answer any four questions: 5×4=20
  - (a) Prove that the set of all feasible solutions of an LPP is a convex set.
  - (b) Sketch the convex polygon spanned by the following points in a two-dimensional Euclidean space. Which of these points are vertices? Express the other as the convex linear combination of the vertices

$$(0,0), (0,1), (1,0), \left(\frac{1}{2}, \frac{1}{4}\right).$$

(c) If  $x_0 \in S$  where S is the set of all FS of the LPP min Z = cx, such that Ax = b,  $x \ge 0$  minimize the objective function Z = cx, then show that  $x_0$  also maximize the objective function  $Z^* = (-c)x$  over S.

(d) Find the dual of the following LPP:  $\min Z_p = x_1 + x_2 + x_3$ 

s.t. 
$$x_1 - 3x_2 + 4x_3 = 5$$
  
 $2x_1 - 3x_2 \le 3$   
 $2x_2 - x_3 \ge 5$   
 $x_1, x_2, x_3 \ge 0$ 

- (e) Prove that the dual of a dual is a primal problem itself.
- (f) Write the characteristics of an LPP in canonical form.
- 4. Answer (a) or (b), (c) or (d), (e) or (f), (g) or (h):
  - (a) Old hens can be bought for Rs. 2 each but young ones cost Rs. 5 each. The old hens lay 3 eggs per week and the young ones 5 eggs per week, each being worth 30 paise. A hen costs Re. 1 per week to feed. If I have only Rs. 80 to spend for hens, how many of each kind shall I buy to give a profit of more than Rs. 6 per week, assuming that I can not house more than 20 hens? Formulate the LPP and solve by graphical method.
  - (b) Find all basic and then all the basic feasible solutions for the equations

$$2x_1 + 6x_2 + 2x_3 + x_4 = 3$$

$$6x_1 + 4x_2 + 4x_3 + 6x_4 = 2$$

and determine the associated general convex combination of the extreme point solutions.

- (c) State and prove the fundamental theorem of LPP.
- (d) Solve by simplex method:

max 
$$Z = 3x_1 + 5x_2 + 4x_3$$
  
s.t.  $2x_1 + 3x_2 \le 8$   
 $3x_1 + 2x_2 + 4x_3 \le 15$   
 $2x_2 + 5x_3 \le 10$   
 $x_1, x_2, x_3 \ge 0$ 

- (e) If in an assignment problem, a constant is added or substracted to every element of a row (or column) of the cost matrix [c<sub>ij</sub>], then prove that an assignment which minimizes the total cost for one matrix, also minimizes the total cost for the other matrix.
- (f) Solve the following transportation problem:

To

From

| DE THE STREET | ALL THE REAL PROPERTY AND ADDRESS OF THE PARTY. |             |   |  |
|---------------|---|-------------|---|--|
| $S_1$         | $S_2$   | $S_3$       | $S_4$   | Supply   |
| 1             | 2   | 1           | 4   | 30   |
|               | 3   | 2           | 1   | 50   |
| -             | 2   | 5           | 9   | 20   |
| 20            | 40  | 30          | 10  | 100  |
|               | 1 3   | 1 2 3 3 4 2 | 1     2     3       1     2     1       3     3     2       4     2     5 | 1     2     3     34       1     2     1     4       3     3     2     1       4     2     5     9 |

(g) For any zero-sum two-persons game where the optimal strategies are not pure and for which A's pay-off matrix is

the optimal strategies are  $(x_1, x_2)$  and  $(y_1, y_2)$  then prove that

$$\frac{x_1}{x_2} = \frac{a_{22} - a_{21}}{a_{11} - a_{12}} \text{ and } \frac{y_1}{y_2} = \frac{a_{22} - a_{12}}{a_{11} - a_{21}} \text{ and }$$
the value of the game to A is given by

$$v = \frac{a_{11}a_{22} - a_{12}a_{21}}{(a_{11} + a_{22}) - (a_{12} + a_{21})}$$

(h) Solve the game whose pay-off matrix is

$$\begin{bmatrix} -1 & -2 & 8 \\ 7 & 5 & -1 \\ 6 & 0 & 12 \end{bmatrix}$$

#### OPTION-B

Paper: MAT-HE-5056

### ( Spherical Trigonometry and Astronomy)

Full Marks: 80

Time: Three hours

## The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions:  $1 \times 10 = 10$ 
  - (a) State one fundamental difference between a spherical triangle and a plane triangle.
  - (b) Define primary circle.
  - (c) Define polar triangle and its primitive triangle.
  - (d) State the third law of Kepler.
  - (e) Explain what is meant by rising and setting of stars.
  - (f) Write any two coordinate systems to locate the position of a heavenly body on the celestial sphere.
  - (g) Define synodic period of a planet.
  - (h) Mention one property of pole of a great circle.

- (i) Just mention how a spherical triangle is formed.
- (j) What is the declination of the pole of the ecliptic?
- 2. Answer the following questions:  $2 \times 5 = 10$ 
  - (a) Prove that section of a sphere by a plane is a circle.
  - (b) Discuss the effect of refraction on sunrise.
  - (c) Drawing a neat diagram, discuss how horizontal coordinates of a heavenly body are measured.
  - (d) Prove that the altitude of the celestial pole at any place is equal to the latitude of that place.
  - (e) Show that right ascension  $\alpha$  and declination  $\delta$  of the sun is always connected by the equation  $\tan \delta = \tan \varepsilon \sin \alpha$ ,  $\varepsilon$  being obliquity of the ecliptic.
- 3. Answer any four of the following:

 $5 \times 4 = 20$ 

(a) Deduce Kepler's laws from Newton's law of gravitation.



- (b) Show that the velocity of a planet in its elliptic orbit is  $v^2 = \mu \left( \frac{2}{r} \frac{1}{a} \right)$  where  $\mu = G(M+m)$  and a is the semi-major axis of the orbit.
- (c) If  $z_1$  and  $z_2$  are the zenith distances of a star on the meridian and the prime vertical respectively, prove that  $\cot \delta = \csc z_1 \sec z_2 \cos z_1$  where  $\delta$  is the star's declination.
- (d) If H be the hour angle of a star of declination  $\delta$  when its azimuth is A and H' when the azimuth is  $(180^{\circ} + A)$ , show that

$$\tan \phi = \frac{\cos \frac{1}{2}(H' + H)}{\cos \frac{1}{2}(H' - H)}$$

- (e) In an equilateral spherical triangle ABC, prove that  $2\cos\frac{a}{2}\sin\frac{A}{2}=1$ .
- (f) If  $\psi$  is the angle which a star makes at rising with the horizon, prove that  $\cos \psi = \sin \phi \sec \delta$ , where the symbols have their usual meanings.

- 4. Answer **any four** questions of the following: 10×4=40
  - (a) If the colatitude is C, prove that  $C = x + \cos^{-1}(\cos x \sec y)$ where  $\tan x = \cot \delta \cos H$  and  $\sin y = \cos \delta \sin H$ , H being the hour angle.
  - (b) In any spherical triangle ABC, prove that  $\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c}$ . Also prove that  $\frac{\sin(A+B)}{\sin C} = \frac{\cos a + \cos b}{1 + \cos c}$
  - (c) Define astronomical refraction and state the laws of refraction. Derive the formula for refraction as  $R = k \tan \xi$ ,  $\xi$  being the apparent zenith distance of a heavenly body. Mention one limitation of this formula.
  - (d) On account of refraction, the circular disc of the sun appears to be an ellipse. Prove it.
  - (e) Derive Kepler's equation in the form  $M = E e \sin E$ , where M and E are respectively mean anomaly and eccentric anomaly.

(f) Assuming the planetary orbits to be circular and coplanar, prove that the sidereal period P and the synodic period S of an inferior planet are related to the earth's periodic time E by

$$\frac{1}{S} = \frac{1}{P} - \frac{1}{E}$$

Calculate the sidereal period (in mean solar days) of a planet whose sidereal period is same as its synodic period.

(g) Prove that, if the fourth and higher powers of e are neglected,

$$E = M + \frac{e \sin M}{1 - e \cos M} - \frac{1}{2} \left( \frac{e \sin M}{1 - e \cos M} \right)^{3}$$

is a solution of Kepler's equation in the form.

(h) Derive the expressions to show the effect of refraction in right ascension and declination.

12

#### OPTION-C

Paper: MAT-HE-5066

( Programming in C)

Full Marks: 60

Time: Three hours

## The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions:  $1 \times 7 = 7$ 
  - (a) Write any two special characters that are used in C.
  - (b) Mention two data types that are used in C language.
  - (c) For x = 2, y = 5, write the output of the C function 'pow (x, y)'.
  - (d) Convert the mathematical expression  $z = e^x + \log y + \sqrt{1+x}$  into C expression.
  - (e) Write the utility of clrscr () function.
  - (f) Write a difference between local variable and global variable.
  - (g) Write the C library function which can evaluate |x|.

- 2. Answer the following questions:  $2 \times 4 = 8$ 
  - (a) Write the difference between 'assignment' and 'equality'.
  - (b) How does 'x + +' differ from '+ + x'?
  - (c) What is a string constant? Give an example.
  - (d) Write four relational operators that are used in C.
- 3. Answer any three parts:  $5\times 3=15$ 
  - (a) Explain artihmetic and logical operators in C with suitable examples.
  - (b) List three header files that are used in C. Also write their utilities. 3+2=5

$$A = 5$$
;  $B = 3$ 

$$A = A + B$$
;

$$B = A - B$$
;

$$A = A - B$$
;

Write the output of A and B from the above program segment in C.

- (c) Write a C program to find the sum of all odd integers between 1 and n.
- (d) Write the general form of do-while loop and explain how it works with the help of a suitable example.

- (e) Write the utility of 'break' and 'continue' statements with the help of suitable examples.
- 4. Why are arrays required in C programming? How are one-dimensional arrays declared and inputs given to array? Explain briefly with example. Write a program to read given numbers and then find the sum of all positive and negative numbers.

  2+3+5=10

#### Or

How are two-dimensional arrays declared? Write a C program to read a 3×3 matrix and print the same as output. Hence write a C program to read a 3×3 matrix, print its transpose and write the determinants of both.

1+4+5=10

- 5. Write a C program for each of the following:
  - (a) To evaluate the function 5  $f(x) = x^2 + 2x 10, x \ge 0$  = |x|, x < 0
  - (b) To find the biggest of three numbers.

5



Explain with example the 'if' statement and nested 'if' statement in C. Write a C program to find the roots of a quadratic equation  $ax^2 + bx + c = 0$ , for all possible values of a,b,c. 5+5=10

6. What is the basic difference between 'Library functions' and 'User-defined functions'? Mention two advantages of using 'User-defined functions'. How are such functions declared and called in a program? Write a C program using function to find the biggest of three numbers.

#### Or

Write a *C* programme that reads a number, obtains a new number by reversing the digits of the given number, and then determine the gcd of the two numbers. To build the programme, use two functions — one to find gcd and another to reverse the digits. 10

