### Total number of printed pages-11

## 3 (Sem-4/CBCS) MAT HC3

#### 2022

#### **MATHEMATICS**

(Honours)

Paper: MAT-HC-4036

(Ring Theory)

Full Marks: 80

Time: Three hours

The figures in the margin indicate full marks for the questions.

1. Answer any ten:

1×10=10

- (a) The set Z of integers under ordinary addition and multiplication is a commutative ring with unity 1. What are the units of Z?
- (b) What is the trivial subring of R?

- (c) What are the elements of  $Z_3[i]$ ?
- (d) Give the definition of zero divisor.
- (e) Give an example of a commutative ring without zero divisors that is not an integral domain.
- (f) What is the characteristic of an integral domain?
- (g) Why is the idea  $\langle x^2 + 1 \rangle$  not prime in  $\mathbb{Z}_2[x]$ ?
- (h) Find all maximal ideals in  $Z_8$ .
- (i) Is the mapping from  $Z_5$  to  $Z_{30}$  given by  $x \rightarrow 6x$  is a ring homomorphism?

- (j) If  $\phi$  is an isomorphism from a ring R onto a ring S, then  $\phi^{-1}$  is an isomorphism from S onto R.

  Write True or False.
- (k) Is the ring 2z isomorphic to the ring 3z?
- (1) Let  $f(x) = x^3 + 2x + 4$  and g(x) = 3x + 2 is  $z_5[x]$ . Determine the quotient and remainder upon dividing f(x) by g(x).
- (m) Why is the polynomial  $3x^5 + 15x^4 20x^3 + 10x + 20$  irreducible over Q?
- (n) Give the definition of Euclidean domain.
- (o) State the second isomorphism theorem for rings.

## Answer any five: $2 \times 5 = 10$

- (a) Define ring. What is the unity of a polynomial ring Z[x]?
- (b) Prove that in a ring R, (-a)(-b) = abfor all  $a, b \in R$ .
- (c) Prove that set S of all matrices of the form  $\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$  with a and b, forms a sub-ring of the ring R of all  $2 \times 2$ matrices having elements as integers.
  - (d) Let R be a ring with unity 1. If 1 has infinite order under addition, then the characteristic of R is 0. If 1 has order n under addition, then prove that the characteristic of R is n.
  - (e) Let  $z/4z = \{0+4z, 1+4z, 2+4z, 3+4z\}.$ Find (2+4z)+(3+4z) and (2+4z)(3+4z)

(f) Let  $R = \left\{ \begin{bmatrix} a & b \\ b & a \end{bmatrix} a, b \in Z \right\}$  and let  $\phi$  be

the mapping defined as  $\begin{bmatrix} a & b \\ b & a \end{bmatrix} \rightarrow a - b$ .

Show that  $\phi$  is a homomorphism.

- (g) Let  $f(x) = 4x^3 + 2x^2 + x + 3$  and  $g(x) = 3x^4 + 3x^3 + 3x^2 + x + 4$ where f(x),  $g(x) \in Z_5[x]$ . Compute f(x)+g(x) and  $f(x)\cdot g(x)$ .
- Prove that in an integral domain, every prime is an irreducible.

# 3. Answer any four:

5×4=20

Define a sub-ring. Prove that a nonempty subset S of a ring R is a subring if S is closed under subtraction and multiplication, that is if a-b and ab are in S whenever a and b are in S. 1+4=5

- (b) Prove that the ring of Gaussian integers  $Z[i] = [a+ib \mid a, b \in Z]$  is an integral domain.
- (c) Let R be a commutative ring with unity and let A be an ideal of R. Then prove that R/A is an integral domain if and only if A is prime.
- (d) If D is an integral domain, then prove that D[x] is an integral domain.
- (e) (i) If R is commutative ring then prove that  $\phi(R)$  is commutative, where  $\phi$  is an isomorphism on R. 3
  - (ii) If the ring R has a unity 1,  $S \neq \{0\}$  and  $\phi: R \rightarrow S$  is onto, then prove that  $\phi(1)$  is the unity of S. 2
- (f) Let  $f(x) \in Z[x]$ . If f(x) is reducible over Q, then prove that it is reducible over Z.

(g) Consider the ring

$$S = \left\{ \begin{bmatrix} a & b \\ -b & a \end{bmatrix} a, b \in Z \right\}.$$
 Show that

 $\phi: \mathbb{C} \to S$  is given by

$$\phi(a+bi) = \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$$
 is a ring

isomorphism.

- (h) Prove that  $Z[i] = \{a + bi | a, b \in Z\}$ , the ring of Gaussian integers is an Euclidean domain.
- 4. Answer any four:

10×4=40

(a) (i) Prove that the set of all continuous real-valued functions of a real variable whose graphs pass through the point (1,0) is a commutative ring without unity under the operation of pointwise addition and multiplication [that is, the operations (f+g)(a) = f(a) + g(a) and  $(f \cdot g)(a) = f(a) \cdot g(a)$ .

- (ii) Prove that if a ring has a unity, it is unique and if a ring element has an inverse, it is unique. 4
- (b) Define a field. Is the set I of all integers a field with respect to ordinary addition and multiplication? Let  $Q\left[\sqrt{2}\right] = \left\{a + b\sqrt{2} \mid a, b \in Q. \text{ Prove that } Q\left[\sqrt{2}\right] \text{ is a field.} \qquad 2+1+7=10$
- (c) (i) Prove that the intersection of any collection of subrings of a ring R is a sub-ring of R.
- (ii) Let R be a commutative ring with unity and let A be an ideal of R. Prove that R/A is a field if A is maximal.

- (d) Define factor ring. Let R be a ring and let A be a subring of R. Prove that the set of co-sets  $\{r+A | r \in R\}$  is a ring under the operation (s+A)+(t+A)=(s+t)+A and (s+A)(t+A)=st+A if and only if A is an ideal of R. 1+5+4=10
- (e) (i) Let  $\phi$  be a ring homomorphism from R to S. Prove that the mapping from  $R/\ker \phi$  to  $\phi(R)$ , given by  $r+\ker \phi \to \phi(r)$  is an isomorphism.
- (ii) Let R be a ring with unity and the characteristic of R is n > 0. Prove that R contains a subring isomorphic to  $Z_n$ . If the characteristic of R is 0, then prove that R contains a sub-ring isomorphic to Z. 3+2=5
  - (f) Let F be a field and let  $p(x) \in F[x]$ . Prove that  $\langle p(x) \rangle$  is a maximal ideal in F[x] if and only if p(x) is irreducible over F.

- (g) Let F be a field and let f(x) and  $g(x) \in F[x]$  with  $g(x) \neq 0$ . Prove that there exists unique polynomials q(x) and r(x) in F[x] such that f(x) = g(x) q(x) + r(x) and either r(x) = 0 or  $degr(x) \langle deg g(x) \rangle$ . With the help of an example verify the division algorithm for F[x].
  - (h) (i) If F is a field, then prove that F[x] is a principal ideal domain. 5
- (ii) Let F be a field and let p(x), a(x),  $b(x) \in F[x]$ . If p(x) is irreducible over F and p(x)|a(x)b(x), then prove that p(x)|a(x) or p(x)|b(x).

(i) Prove that every principal ideal domain is a unique factorization domain.

- (j) (i) Prove that every Euclidean domain is a principal ideal domain. 5
  - (ii) Show that the ring  $Z\left[\sqrt{-5}\right] = \left\{a + b\sqrt{-5} \mid ab \in Z\right\}$  is an integral domain but not a unique factorization domain. 5

